

60. You are given the following information about six coins:

Coin	Probability of Heads
1 – 4	0.50
5	0.25
6	0.75

A coin is selected at random and then flipped repeatedly. X_i denotes the outcome of the i th flip, where “1” indicates heads and “0” indicates tails. The following sequence is obtained:

$$S = \{X_1, X_2, X_3, X_4\} = \{1, 1, 0, 1\}$$

Calculate $E(X_5 | S)$ using Bayesian analysis.

- (A) 0.52
- (B) 0.54
- (C) 0.56
- (D) 0.59
- (E) 0.63

61. You observe the following five ground-up claims from a data set that is truncated from below at 100:

$d=100$

125 150 165 175 250

You fit a ground-up exponential distribution using maximum likelihood estimation.

Calculate the mean of the fitted distribution.

- (A) 73
- (B) 100
- (C) 125
- (D) 156
- (E) 173



→: The log-likelihood f'tion is:

$$l(\theta) = \sum_{j=1}^n \ln(f_X(x_j; \theta)) - n \cdot \ln(S_X(d; \theta))$$

We have $X \sim \text{Exponential}(\theta)$

$$l(\theta) = \sum_{j=1}^n \ln\left(\frac{1}{\theta} \cdot e^{-\frac{x_j}{\theta}}\right) + n \cdot \cancel{\ln\left(e^{-\frac{d}{\theta}}\right)}$$

$$l(\theta) = \sum_{j=1}^n \left(-\ln(\theta) + \cancel{\ln\left(e^{-\frac{x_j}{\theta}}\right)} \right) + n \cdot \frac{d}{\theta}$$

$$l(\theta) = -n \cdot \ln(\theta) + \sum_{j=1}^n \left(-\frac{x_j}{\theta} \right) + n \cdot \frac{d}{\theta}$$

$$l(\theta) = -n \cdot \ln(\theta) - \frac{1}{\theta} \sum_{j=1}^n x_j + n \cdot \frac{d}{\theta}$$

We differentiate

$$l'(\theta) = -n \cdot \frac{1}{\theta} + (+1) \cdot \frac{1}{\theta^2} \sum_{j=1}^n x_j + n(-1) \cdot \frac{d}{\theta^2} = 0$$

$$\underbrace{\left(\frac{1}{\theta^2} \left(-n \cdot \theta + \sum_{j=1}^n x_j - n \cdot d \right) \right)}_{=0} = 0$$

$$\hat{\theta}_{MLE} = \left(\frac{1}{n} \sum_{j=1}^n x_j \right) - d = \bar{x} - d$$

Task:
Repeat for

$X \sim \text{Gamma}(\alpha=2, \theta)$

In this problem:

$$\hat{\theta}_{MLE} = \frac{1}{5}(125 + 150 + 165 + 175 + 250) - 10$$

$$\hat{\theta}_{MLE} = 73$$



152. You are given:

- (i) A sample of losses is:

600 700 900

- (ii) No information is available about losses of 500 or less. **truncation** **$d=500$**

- (iii) Losses are assumed to follow an exponential distribution with mean θ .

Calculate the maximum likelihood estimate of θ .

(A) 233

(B) 400

(C) 500

 (D) 733

(E) 1233

→ :

$$\hat{\theta}_{MLE} = \bar{x} - d$$

$$\hat{\theta}_{MLE} = \frac{1}{3}(600+700+900) - 500$$

$$\hat{\theta}_{MLE} = \frac{2200}{3} - 500 = 733.\bar{3} - 500$$

$$\boxed{\hat{\theta}_{MLE} \approx 233}$$

□

153. DELETED

261. DELETED

262. You are given:

- (i) At time 4 hours, there are 5 working light bulbs.) truncated @ 4
- (ii) The 5 bulbs are observed for p more hours. complete data
- (iii) Three light bulbs burn out at times 5, 9, and 13 hours, while the remaining light bulbs are still working at time $4 + p$ hours. censoring
- (iv) The distribution of failure times is uniform on $(0, \omega)$.
- (v) The maximum likelihood estimate of ω is 29

Calculate p .

- (A) Less than 10
- (B) At least 10, but less than 12
- (C) At least 12, but less than 14
- (D) At least 14, but less than 16
- (E) At least 16

$$X \sim U(0, \omega)$$

$$\text{pdf: } f_X(x; \omega) = \frac{1}{\omega}$$

for $x \in (0, \omega)$

survival f'tion:

$$S_X(x; \omega) = \frac{\omega - x}{\omega}$$

for $x \in (0, \omega)$

The Likelihood f'tion:

$$L(\omega) = \frac{\left(\frac{1}{\omega}\right)^3 \cdot (S_X(4+p; \omega))^2}{(S_X(4; \omega))^5}$$

$$L(\omega) = \frac{\left(\frac{1}{\omega}\right)^3 \cdot \left(\frac{\omega - (4+p)}{\omega}\right)^2}{\left(\frac{\omega - 4}{\omega}\right)^5}$$

$$L(\omega) = \frac{(\omega - 4 - p)^2}{(\omega - 4)^5}$$

The log-likelihood:

$$l(w) = 2 \cdot \ln(w-4-p) - 5 \cdot \ln(w-4)$$

$$l'(w) = \frac{2}{w-4-p} - \frac{5}{w-4}$$

We are given that $\hat{w}_{MLE} = 29 \Rightarrow l'(29) = 0$

$$\frac{2}{29-4-p} = \frac{5}{29-4} = \frac{1}{5}$$

$$10 = 25-p \Rightarrow \boxed{p=15}$$

□

MLE for Discrete Distributions

MLE: Bernoulli.

$$\left(\begin{array}{l} X \sim \text{Bernoulli}(q) \\ \quad \quad \quad \uparrow \text{the probab. of success} \\ \text{Support}(X) = \{0, 1\} \\ X \sim \begin{cases} 0 & \text{w/ probab. } 1-q \\ 1 & \text{w/ probab. } q \end{cases} \end{array} \right)$$

Let $x_1, x_2, \dots, x_n$ be the observations from Bernoulli(q).

They will all be 0 or 1.

We write the pmf as:

$$f(x; q) = \begin{cases} q & \text{if } x=1 \\ 1-q & \text{if } x=0 \end{cases}$$

$$\boxed{f(x; q) = q^x \cdot (1-q)^{1-x}}$$