

UNIVERSITY OF TEXAS AT AUSTIN

Problem Set # 20

The t -procedure: two means20.1. Two-sample t .

Problem 20.1. An instructor is teaching two sections of the same basic statistics course. The instructor is giving the same exams, homework assignments, and quizzes in both sections. Which t -procedure should be used to determine if there is a difference in the academic performance between the two course sections?

- (a) One-sample t -test.
- (b) Matched-pairs t -procedure.
- (c) Two-sample t -test.
- (d) None of the above.

Problem 20.2. This is an excerpt from findings of an educational study:

A study was done to determine whether there is a difference in the amount of time (in hours) that graduate students versus undergraduate students spend on the Internet per day. Five undergrads and five grad students were polled.

- (i) Is the alternative hypothesis one-sided or two-sided?
- (ii) A t -score for the data gathered was calculated to be 1.6664. Would you say that there is a significant difference in the amount of time that graduate and undergraduate students spend on the Internet?

$$p\text{-value} = 2 * pt(-1.6664, df=4) = 0.1709623$$

No!

Problem 20.3. (5 points)

There is a dispute about salaries of male versus female elves. The North Polar Bear collected the following data:

- the total number of male elves is 121;
- the total number of female elves is 100;
- the average salary of a male elf is 10,000 candy canes;
- the average salary of a female elf is 12,000 candy canes;
- the sample standard deviation of the salaries of male elves is 50;
- the sample standard deviation of the salaries of female elves is 132.

Assume independence between the salaries of individual elves.

Let μ_m denote the population mean for the distribution of the male elves' salaries and let μ_f denote the population mean for the distribution of the female elves' salaries. We wish to test:

$$H_0 : \mu_m = \mu_f \text{ vs. } H_a : \mu_m \neq \mu_f.$$

What is the p -value associated with our data?

- a.: About 0.
- b.: About 0.01.
- c.: About 0.025
- d.: About 0.04.
- e.: None of the above.

→: Two-sided t -test

$$t = \frac{\bar{x}_m - \bar{x}_f}{\sqrt{\frac{s_m^2}{n_m} + \frac{s_f^2}{n_f}}} = \frac{10000 - 12000}{\sqrt{\frac{50^2}{121} + \frac{132^2}{100}}} \Rightarrow$$

$$t = -143.26 \Rightarrow \text{the } p\text{-value is negligible}$$

$$df = \min(n_1, n_2) - 1 = 99$$

Problem 20.4. Let the population distributions be normal with unknown parameters. Assume that sample data, based on two independent samples of size 25, give us $\bar{x}_1 = 505$, $\bar{x}_2 = 515$, $s_1 = 23$, and $s_2 = 28$.

- (i) What is a 95% confidence interval (use the conservative value for the degrees of freedom) for the difference between the two population means?
- (ii) Based on the confidence interval, we can conclude at the 5% significance level that there is no difference between the two population means. *True or false?*
- (iii) The margin of error for the difference between the two sample means would be smaller if we were to take larger samples. *True or false?*
- (iv) If a 99% confidence interval were calculated instead of the 95% interval, it would include more values for the difference between the two population means. *True or false?*

20.2. Pooled t (optional content from Subsection 7.3.4).

Problem 20.5. The pooled two-sample t -procedure can be used when ...

- (a) you can assume the two populations have equal variances
- (b) you can assume the two populations have equal means
- (c) the sample sizes are equal
- (a) None of the above

Problem 20.6. Let n_1 and n_2 denote the sample sizes of each group. The pooled two-sample t -procedure is based how many degrees of freedom?

- (a) $n_1 + n_2 + 2$
- (b) $n_1 + n_2 - 2$
- (c) $n_1 + n_2 - 1$
- (d) $n_1 + n_2$
- (e) None of the above.

Problem 20.7. A study was done to determine if students learn better in an online basic statistics class versus a traditional face-to-face (f2f) course. A random sample of 12 students in an online course and 15 students in an f2f course was taken.

- (i) Let μ_{new} denote the population mean score for the online statistics class and let μ_{old} denote the population mean score for the face-to-face statistics class. What are the hypotheses being tested?
- (ii) We decide it is appropriate to use the pooled t -procedure. What is the number of degrees of freedom you are going to use?

i.

$$\boxed{\text{pt. estimate} \pm \text{margin of error}}$$

- pt. estimate for $\mu_1 - \mu_2$: $\bar{x}_1 - \bar{x}_2 = 505 - 515 = -10$

- margin of error = $t^* \cdot \text{std error}$

- $\text{std error} = \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}} = \sqrt{\frac{23^2}{25} + \frac{28^2}{25}} =$
 $= \frac{\sqrt{23^2 + 28^2}}{5} = 7.247$

- $t^* = ?$ $df = 25 - 1 = 24$ $\left. \begin{array}{l} \text{upper tail probab.} = 0.025 \end{array} \right\} \Rightarrow t^* = 2.064$
 $\uparrow$
t-tables

$$\Rightarrow \underline{\mu_1 - \mu_2 = -10 \pm 2.064(7.247) = -10 \pm 14.96}$$