

M358K: October 9th, 2023.

Confidence Intervals [The Normal Case].

We are still in the normal model.

The same logic will apply to other models as well.

Let $X_1, X_2, \dots, X_n$ be a normal random sample, i.e.,
 $\{X_i, i=1..n\}$ are all independent, and
 $X_i \sim \text{Normal}(\text{mean} = \mu, \text{sd} = \sigma)$

?"


We know exactly the distribution of the sample mean:

$$\bar{X}_n \sim \text{Normal}(\text{mean} = \mu, \text{sd} = \frac{\sigma}{\sqrt{n}})$$

?"

For now: assume that σ is known.

we know that $\bar{X}_n$ is a "good" estimator for the population mean μ



Q: How **CONFIDENT** are we about the value that we get?
What does "confidence" even mean?

Let C be a large probability, i.e., a confidence level.

Say $C = 0.95, 0.90, 0.99, 0.80$



Look @

$$\mathbb{P} \left[|\bar{X} - \mu| < (?) \right] = C$$

$$\mathbb{P} \left[-(?) < \bar{X} - \mu < (?) \right] = C$$

$$\mathbb{P} \left[\frac{-(?)}{\frac{\sigma}{\sqrt{n}}} < \frac{\bar{X} - \mu}{\frac{\sigma}{\sqrt{n}}} < \frac{(?)}{\frac{\sigma}{\sqrt{n}}} \right] = C$$

$-z^* \quad \bar{Z} \quad z^* \quad \sim N(0,1)$



$$\frac{1-C}{2} + C = \frac{1+C}{2}$$

$$z^* = \Phi^{-1} \left(\frac{1+C}{2} \right)$$

$$\longleftrightarrow \text{qnorm}((1+C)/2)$$

z^* is the critical value of $N(0,1)$ such that

$$\mathbb{P}[-z^* < Z < z^*] = C$$



$$\Rightarrow z^* = \frac{(?)}{\frac{\sigma}{\sqrt{n}}} \Rightarrow$$

$$(?) = z^* \frac{\sigma}{\sqrt{n}}$$

$$\textcircled{\checkmark} \Rightarrow \mathbb{P}\left[-z^* \cdot \frac{\sigma}{\sqrt{n}} < \bar{X} - \mu < z^* \cdot \frac{\sigma}{\sqrt{n}}\right] = C$$

$$\Rightarrow \mathbb{P}\left[\bar{X} - z^* \cdot \frac{\sigma}{\sqrt{n}} < \mu < \bar{X} + z^* \cdot \frac{\sigma}{\sqrt{n}}\right] = C$$

Random Interval

which we call our CONFIDENCE INTERVAL

- Every time that you collect a sample and construct a confidence interval, you will obtain a different interval.
- The confidence interval will contain the mean parameter μ with probability C , and w/ probability $1-C$, it will NOT.

In Practice:

For a particular data set: $x_1, x_2, \dots, x_n$,

we calculate $\bar{x} = \frac{1}{n}(x_1 + \dots + x_n)$ sample average

Then, we provide C -confidence interval:

$$\left(\bar{x} - z^* \cdot \frac{\sigma}{\sqrt{n}}, \bar{x} + z^* \cdot \frac{\sigma}{\sqrt{n}}\right).$$

We usually write:

$$\mu = \bar{x} \pm z^* \cdot \frac{\sigma}{\sqrt{n}}$$

In words: "We are C -confident that the mean μ is within the interval...."

$z^* \cdot \frac{\sigma}{\sqrt{n}}$ is called the margin of error in this particular case

More generally, the margin of error is the radius of the confidence interval.

