

M362K: April 22nd, 2024.

Section 4.2.

The Exponential Distribution.

A r.v. T is said to have the exponential distribution with rate λ , where λ is a positive parameter, if its pdf is of this form

$$f_T(t) = \lambda e^{-\lambda \cdot t} \quad t \geq 0$$

The cumulative dist'n f'n of T is, then,

$$\begin{aligned} F_T(t) &= \int_0^t f_T(u) du = \int_0^t \lambda e^{-\lambda \cdot u} du = \\ &= \lambda \cdot \int_0^t e^{-\lambda u} du = \cancel{\lambda} \cdot \left(-\frac{1}{\cancel{\lambda}} \right) e^{-\lambda u} \Big|_{u=0}^t \\ &= \underline{1 - e^{-\lambda \cdot t}} \quad \text{for } t \geq 0 \end{aligned}$$

Def'n. For a r.v. X w/ cdf F_X , its survival function is defined as

$$S_X(x) = \mathbb{P}[X > x] = 1 - F_X(x) \quad \text{for all } x \in \mathbb{R}$$

Properties.

• W/out proof:

$$\underline{\mathbb{E}[T] = \text{SD}[T] = \frac{1}{\lambda}}$$

• $\underline{S_T(t) = e^{-\lambda t}}$

Problem. The amt of time, in hrs, that a computer functions before breaking down is a continuous r.v. w/ pdf

$$f_T(t) = \begin{cases} \lambda e^{-\frac{t}{100}} & \text{for } t \geq 0 \\ 0 & x < 0 \end{cases}$$

(i) What is the probability that the computer will function for less than 100 hrs?

$$\rightarrow: T \sim \text{Exponential} \left(\lambda = \frac{1}{100} \right) \Rightarrow \lambda = \frac{1}{100}$$

$$\mathbb{P}[T \leq 100] = F_T(100) = 1 - e^{-\frac{100}{100}} = 1 - e^{-1} = 0.632$$

(ii) What is the probability that the computer will function between 50 and 150 hrs before breaking down?

$$\begin{aligned}\rightarrow: \mathbb{P}[50 < T < 150] &= F_T(150) - F_T(50) \\ &= 1 - e^{-\frac{150}{100}} - \left(1 - e^{-\frac{50}{100}}\right) = \\ &= e^{-\frac{50}{100}} - e^{-\frac{150}{100}} = \underline{S_T(50) - S_T(150)} \\ &= e^{-0.5} - e^{-1.5}\end{aligned}$$

□

Half-Life.

Let m denote the half-life of $T \sim \text{Exp}(\lambda)$, i.e., let m satisfy:

$$\mathbb{P}[T \leq m] = \frac{1}{2} = \mathbb{P}[T \geq m] \quad (T \text{ continuous})$$

$$\frac{1}{2} = S_T(m)$$

$$\frac{1}{2} = e^{-\lambda \cdot m}$$

$$+\ln(2) = \ln\left(\frac{1}{2}\right) = -\lambda \cdot m$$

$$m = \frac{\ln(2)}{\lambda}$$

In other words
 $\frac{\ln(2)}{\lambda}$ is the median
of $\text{Exp}(\lambda)$.

Note: In the previous problem,

- the half-life would be $m = \frac{\ln(2)}{\frac{1}{100}} = 100 \ln(2)$
- the mean would be $\frac{1}{\lambda} = 100$

Memoryless Property .

Let $T \sim \text{Exp}(\lambda)$. Let $s, t \geq 0$:

$$\underline{\mathbb{P}[T > s+t \mid T > t]} = \frac{\mathbb{P}[\overset{s}{T > s+t}, \overset{s}{T > t}]}{\mathbb{P}[T > t]} = \frac{\mathbb{P}[T > s+t]}{\mathbb{P}[T > t]} =$$

by def'n
of conditional
probability

$$= \frac{e^{-\lambda(s+t)}}{e^{-\lambda t}} = e^{-\lambda s} = S_T(s) = \underline{\mathbb{P}[T > s]}$$

survival
f'n

28. The number of days that elapse between the beginning of a calendar year and the moment a high-risk driver is involved in an accident is exponentially distributed. An insurance company expects that 30% of high-risk drivers will be involved in an accident during the first 50 days of a calendar year.

Calculate the portion of high-risk drivers are expected to be involved in an accident during the first 80 days of a calendar year.

- (A) 0.15
(B) 0.34
(C) 0.43
(D) 0.57
(E) 0.66

$T \dots \# \text{ of days before a h.r driver has an accident}$
 $P[T \leq 50] = 0.3 \Rightarrow P[T > 50] = 0.7 \Rightarrow e^{-\lambda \cdot 50} = 0.7$
 $P[T \leq 80] = 1 - e^{-80\lambda} = 1 - (e^{-10\lambda})^8 = 1 - (0.7)^{8/5}$
 $e^{-10\lambda} = (0.7)^{1/5}$

29. An actuary has discovered that policyholders are three times as likely to file two claims as to file four claims.

The number of claims filed has a Poisson distribution.

Calculate the variance of the number of claims filed.

- (A) $\frac{1}{\sqrt{3}}$
(B) 1
(C) $\sqrt{2}$
(D) 2
(E) 4

30. A company establishes a fund of 120 from which it wants to pay an amount, C , to any of its 20 employees who achieve a high performance level during the coming year. Each employee has a 2% chance of achieving a high performance level during the coming year. The events of different employees achieving a high performance level during the coming year are mutually independent.

Calculate the maximum value of C for which the probability is less than 1% that the fund will be inadequate to cover all payments for high performance.

- (A) 24
(B) 30
(C) 40
(D) 60
(E) 120

266. Claim amounts are independent random variables with probability density function

$$f(x) = \begin{cases} \frac{10}{x^2}, & \text{for } x > 10 \\ 0, & \text{otherwise.} \end{cases}$$

Calculate the probability that the largest of three randomly selected claims is less than 25.

- (A) $\frac{8}{125}$
- (B) $\frac{12}{125}$
- (C) $\frac{27}{125}$
- (D) $\frac{2}{5}$
- (E) $\frac{3}{5}$

267. The lifetime of a certain electronic device has an exponential distribution with mean 0.50.

Calculate the probability that the lifetime of the device is greater than 0.70, given that it is greater than 0.40.

- (A) 0.203
- (B) 0.247
- (C) 0.449
- (D) 0.549
- (E) 0.861

→: lifetime $T \sim \text{Exp}(\lambda=2)$

$$\begin{aligned} \mathbb{P}[T > 0.7 \mid T > 0.4] &= \mathbb{P}[T > 0.3] = S_T(0.3) \\ &\stackrel{\text{memoryless property}}{=} e^{-2(0.3)} \\ &= e^{-0.6} \end{aligned}$$



Example. 4.5.3. Minimum of Independent Exponential r.v.s.

Let $X_1, X_2, \dots, X_n$ be independent r.v.s

w/ $X_i \sim \text{Exp}(\lambda_i)$ for $i=1, \dots, n$.

Define: $X_{\min} = \min(X_1, X_2, \dots, X_n)$.

Let's find the cdf of $X_{\min}$.

$$\begin{aligned} \text{For every } x > 0: \quad F_{X_{\min}}(x) &= \mathbb{P}[X_{\min} \leq x] = 1 - \mathbb{P}[X_{\min} > x] \\ &= 1 - \mathbb{P}[X_1 > x, X_2 > x, \dots, X_n > x] \\ &\quad \text{independence} \\ &= 1 - S_{X_1}(x) \cdot S_{X_2}(x) \cdot \dots \cdot S_{X_n}(x) \\ &= 1 - e^{-\lambda_1 x} \cdot e^{-\lambda_2 x} \cdot \dots \cdot e^{-\lambda_n x} \\ &= 1 - e^{-(\lambda_1 + \lambda_2 + \dots + \lambda_n) \cdot x} \end{aligned}$$

$X_{\min} \sim \text{Exp}(\lambda = \lambda_1 + \dots + \lambda_n)$

248. Let X be a random variable that takes on the values -1 , 0 , and 1 with equal probabilities.

Let $Y = X^2$.

Which of the following is true?

- (A) $\text{Cov}(X, Y) > 0$; the random variables X and Y are dependent.
- (B) $\text{Cov}(X, Y) > 0$; the random variables X and Y are independent.
- (C) $\text{Cov}(X, Y) = 0$; the random variables X and Y are dependent.
- (D) $\text{Cov}(X, Y) = 0$; the random variables X and Y are independent.
- (E) $\text{Cov}(X, Y) < 0$; the random variables X and Y are dependent.

249. Losses follow an exponential distribution with mean 1. Two independent losses are observed.

Calculate the expected value of the smaller loss.

- (A) 0.25
- (B) 0.50
- (C) 0.75
- (D) 1.00
- (E) 1.50

250. A delivery service owns two cars that consume 15 and 30 miles per gallon. Fuel costs 3 per gallon. On any given business day, each car travels a number of miles that is independent of the other and is normally distributed with mean 25 miles and standard deviation 3 miles.

Calculate the probability that on any given business day, the total fuel cost to the delivery service will be less than 7.

- (A) 0.13
- (B) 0.23
- (C) 0.29
- (D) 0.38
- (E) 0.47