

M362K: February 16th, 2024.

Geometric Distribution [revisited].

Start w/ a sequence of independent, identically distributed Bernoulli trials.
 the same probability p of success in a single trial.

Repeat the trials until the first success.
Count how many trials it took (including the success).

Outcome Space:

$$\Omega = \mathbb{N} = \{1, 2, 3, \dots\}$$

Distribution:

$$p_k := \mathbb{P}[\{k\}] = (1-p)^{k-1} \cdot p \text{ for all } k \in \mathbb{N}$$

Example. Consider a house in a 100-yr flood plane, then this can be interpreted as $p = \frac{1}{100}$ being the chance of a flood in a single year.

Assume independent years.

$$\mathbb{P}[\text{no flood in next 30 years}] = \left(\frac{99}{100}\right)^{30} = 0.7397$$

UNIVERSITY OF TEXAS AT AUSTIN

Problem Set # 2

Sequences of events.

Problem 2.1. Roger is playing darts. His throws are all mutually independent, and he has a probability 0.3 of hitting the bull's eye in any single throw. How many darts should Roger throw so that there is at least an 80% probability of hitting bull's eye at least once?

→: $P[\text{no bulls eye in } n \text{ throws}] = (0.7)^n$

$$1 - (0.7)^n \geq 0.8$$

$$(0.7)^n \leq 0.2$$

ln |

$$n \cdot \ln(0.7) \leq \ln(0.2) \quad /: \ln(0.7) < 0$$

$$n \geq \frac{\ln(0.2)}{\ln(0.7)} = 4.51 \quad \Rightarrow \quad \boxed{n \geq 5}$$



Problem 2.2. Audrey and Evie take turns tossing a fair coin with Audrey having the first turn. Whoever gets Heads first wins the game.

- (i) What's the probability that Evie wins the game?
 (ii) Is it possible to weight the coin so that the game is fair, i.e., with what probability p should Heads appear so that Audrey and Evie are equally likely to win the game?

p_E ... probab. that Evie wins

$$p_E = \text{TP}[\text{it takes an even \# of trials}]$$

$$= p_2 + p_4 + \dots + p_{2k} + \dots$$

$$= \underbrace{(1-p) \cdot p}_a + (1-p)^3 \cdot p + \dots + (1-p)^{2k-1} \cdot p + \dots$$

$$= (1-p) \cdot p (1 + (1-p)^2 + (1-p)^4 + \dots)$$

$$= \frac{(1-p)p}{1-(1-p)^2} = \frac{(1-p)p}{(1-1+p)(1+1-p)} = \frac{1-p}{2-p}$$

For $p = \frac{1}{2}$:

$$p_E = \frac{1}{3}$$

(ii) Try to solve:

$$\frac{1-p}{2-p} = \frac{1}{2}$$

$$\cancel{2} - 2p = \cancel{2} - p$$


$p=0$ means they play forever

Method I.

Method II:

p_E ... probab. that Evie wins

p_A ... probab. that Audrey wins



$$p_E = (1-p) p_A$$

$$p_E + p_A = 1$$

$$(1-p) p_A + p_A = 1$$

$$(2-p) p_A = 1$$

$$p_A = \frac{1}{2-p}$$

$\Rightarrow$

$$p_E = \frac{1-p}{2-p}$$



Problem 2.3. Source: Problem #1.5.6 from Pitman.

Suppose you roll a fair six-sided die repeatedly until the first time you roll a number that you have rolled before. Let p_r denote the probability that you roll the die exactly r times.

- (i) Without calculation, write down the value of $p_1 + p_2 + \dots + p_{10}$. Explain. 1
- (ii) For each $r = 1, 2, \dots$ calculate the probability p_r . ✓
- (iii) Verify arithmetically that your response to (i) was correct.


$p_1 = 0$
 $p_2 = \frac{1}{6}$
 $p_3 = \frac{5}{6} \cdot \frac{2}{6}$
 $p_4 = \frac{5}{6} \cdot \frac{4}{6} \cdot \frac{3}{6}$
 $p_5 = \frac{5}{6} \cdot \frac{4}{6} \cdot \frac{3}{6} \cdot \frac{2}{6}$
 $p_6 = \frac{5}{6} \cdot \frac{4}{6} \cdot \frac{3}{6} \cdot \frac{2}{6} \cdot \frac{1}{6}$
 $p_7 = \frac{5}{6} \cdot \frac{4}{6} \cdot \frac{3}{6} \cdot \frac{2}{6} \cdot \frac{1}{6} \cdot 1$

$p_r = 0$ for $r \geq 8$



Section 2.1.


Binomial Distribution.



w/ probab. p^2

w/ probab. $2p(1-p)$

w/ probab. $(1-p)^2$



w/ probab. $\binom{n}{k} p^k (1-p)^{n-k}$

Outcome Space: $\Omega = \{0, 1, 2, \dots, n-1, n\}$

Distribution:

$$p_k = \binom{n}{k} p^k (1-p)^{n-k} \quad k \in \Omega$$

We write:

Binomial(n, p) or $b(n, p)$