

M362K: January 19th, 2024.

Reading Assignment: Appendices 1-4.

Section 1.1. Inspiration

"Equally likely outcomes"

(aka "picked/chosen @ random")

Ω ... set of all possible elementary outcomes ω

Note: $\omega \in \Omega$

Start w/ finite for simplicity.

e.g., cointoss $\Omega = \{H, T\}$; $\#(\Omega) = 2$;

roll of a die $\Omega = \{1, 2, 3, 4, 5, 6\}$; $\#(\Omega) = 6$;

rock-paper-scissors $\Omega = \{R, P, S\} \times \{R, P, S\}$
 $= \{(R, R), (R, P), (R, S),$
 $(P, R), (P, P), (P, S),$
 $(S, R), (S, P), (S, S)\}$; $\#(\Omega) = 9 = 3 \times 3$

fastcasual restaurant:

$\Omega = \{\text{nice, pasta}\} \times \{\text{tofu, salmon, chicken}\} \times \{\text{carrots, peas, beans}\}$
 $\#(\Omega) = 2 \times 3 \times 3 = 18$

Any "nice" subset of Ω is called an **event**;

e.g., a prime number on the die $A = \{2, 3, 5\}$; $\#(A) = 3$

the first player wins $B = \{(R, S), (P, R), (S, P)\}$;

vegetarian meals: $E = \{\text{nice, pasta}\} \times \{\text{tofu}\} \times \{\text{carrots, peas, beans}\}$ $\#(E) = 6$

meals w/ beans: $F = \{\text{nice, pasta}\} \times \{\text{tofu, salmon, chicken}\} \times \{\text{beans}\}$

veg meals w/ beans: $G = \{\text{nice, pasta}\} \times \{\text{tofu}\} \times \{\text{beans}\}$

$$= E \cap F$$

If all outcomes are equally likely, that's the same as if we said they have the same probability/chance/likelihood.

It's natural to have $\mathbb{P}[\Omega] = 1$.

So,

$$\mathbb{P}[\underbrace{\{\omega\}}_{\text{singleton}}] = \frac{1}{\#(\Omega)} \text{ for all } \omega$$

In fact, for every event $E \subseteq \Omega$, it's natural to set

$$\mathbb{P}[E] = \frac{\#(E)}{\#(\Omega)}$$

e.g., $\mathbb{P}[\text{prime number on die}] = \frac{3}{6} = \frac{1}{2}$

$$\mathbb{P}[\text{veg. meal}] = \frac{\cancel{2} \times 1 \times \cancel{3}}{\cancel{2} \times 3 \times \cancel{3}} = \frac{1}{3}$$

Example. [Example 3]

You roll two n -sided dice w/ sides numbered $1, 2, 3, \dots, n$.
(w/ $n \geq 4$).

Q: What's the probability that the sum is 5?

$$\rightarrow: \Omega = \{(i, j) : 1 \leq i, j \leq n\}$$

$$\#(\Omega) = n \times n = n^2$$


$$H = \{\text{sum is 5}\} = \{(1, 4), (2, 3), (3, 2), (4, 1)\}$$

$$\mathbb{P}[H] = \frac{\#(H)}{\#(\Omega)} = \frac{4}{n^2}$$

□

Q: What's the probability that the number on the second die is strictly greater?

→:



$D = \{\text{same on both dice}\}$

$E = \{\text{second greater}\}$

$C = \{\text{first greater}\}$

$$\#(D) = n$$

symmetry $\#(E) = \#(C)$

$$\#(\Omega) = \#(E) + \#(D) + \#(C) = 2\#(E) + \#(D) \Rightarrow \#(E) = \frac{\#(\Omega) - \#(D)}{2} = \frac{n^2 - n}{2}$$

$$\mathbb{P}[E] = \frac{\#(E)}{\#(\Omega)} = \frac{\frac{1}{2}(n^2 - n)}{n^2} = \frac{1}{2} \left(1 - \frac{1}{n}\right) \xrightarrow{n \rightarrow \infty} \frac{1}{2}$$

$\downarrow n \rightarrow \infty$
0

Problem. Balls in an urn: Y, B, R, G.

Twice as many Y as B. $\Rightarrow y = 2b = 8g$
 Twice as many B as R. $\Rightarrow b = 2r = 4g$
 Twice as many R as G. $\Rightarrow r = 2g$

$\mathbb{P}[G] = ?$

$\rightarrow y, b, r, g$

$$1 = g + 2g + 4g + 8g$$

$$g = \frac{1}{15}$$

□