

M362K: March 4th, 2024.

Section 3.1.

Random Variables

... on a finite Ω (for now).

Example. Consider an experiment consisting of 3 independent tosses of a fair coin.
What is the outcome space Ω corresponding to this experiment?

$$\Omega = \{HHH, THH, HTH, HHT, TTH, THT, HTT, TTT\}$$

Q: How do we model the number of Hs in the first two coin tosses?

$$\Omega = \{HHH, THH, HTH, HHT, TTH, THT, HTT, TTT\}$$

$$\begin{array}{cccccccc} \text{X} \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \\ 2 & 1 & 1 & 2 & 0 & 1 & 1 & 0 \end{array}$$

← Now we have a function from Ω into $\mathbb{R}$.

Def'n. Let Ω be a finite outcome space.

Any function

$$\text{X}: \Omega \rightarrow \mathbb{R}$$

is called a **RANDOM VARIABLE.**

Note: By convention, random variables are usually denoted by the capital letters of the latin alphabet.

$$\Omega = \{\omega^1, \omega^2, \dots, \omega^N\}$$
$$\begin{array}{cccc} \text{X} \downarrow & \downarrow & \searrow & \swarrow \\ & & & \end{array}$$

Support(X) = $\{x_1, x_2, \dots, x_m\} \subseteq \mathbb{R}$ is every possible value X can take

Def'n. The **probability (mass) function (pmf)** of X is defined as follows:

$$p_X(k) := \mathbb{P}[X = x_k] \quad \text{for all } k=1, \dots, m$$

Example [cont'd].

$$\text{Support}(X) = \{0, 1, 2\}$$

It's usually convenient to place the **pmf** in a table:

k	0	1	2
$p_X(k)$	$\frac{1}{4}$	$\frac{1}{2}$	$\frac{1}{4}$

In general:

$$\sum_{k=1}^m p_X(x_k) = 1$$

Example. Bernoulli distribution w/ success probability p .

$$\Omega = \{0, 1\}$$

$$\begin{array}{c} \times \\ \downarrow \downarrow \\ \text{Support}(X) = \{0, 1\} \end{array}$$

The pmf of X :

$$p_X(0) = \mathbb{P}[X=0] = 1-p$$

$$p_X(1) = \mathbb{P}[X=1] = p$$

We write:

$$X \sim \text{Bernoulli}(p)$$


$$\text{OR} \\ X \sim b(p)$$

We say: X has the Bernoulli dist'n w/ parameter p

OR X is Bernoulli dist'd w/ parameter p .

Example.

Consider an event A on an outcome space Ω .



We define an indicator random variable of the event A .

$$\omega \in \Omega: I_A(\omega) = \begin{cases} 1 & \text{if } \omega \in A \\ 0 & \text{if } \omega \notin A \end{cases}$$

$$\text{Support}(I_A) = \{0, 1\}$$

The pmf of I_A :

k	0	1
$p_{I_A}(k)$	$1 - P[A]$	$P[A]$

In other words:

$$I_A \sim \text{Bernoulli}(p = P[A])$$

Example. Binomial w/ n trials and success probability p .

It's sufficient: $\Omega = \{0, 1, \dots, n-1, n\}$

Support(X) = $\{0, 1, \dots, n-1, n\}$

Its pmf is:

for all $k = 0, \dots, n$:

$$p_X(k) = \binom{n}{k} p^k (1-p)^{n-k}$$

We write: $X \sim \text{Binomial}(n, p)$ or $X \sim b(n, p)$

We say: X is binomial w/ parameters n and p .

Example. Consider rolling a fair ordinary die.

$$\Omega = \{1, 2, 3, 4, 5, 6\}$$

Let the random variable X be the value on the upturned face.

$$\begin{aligned} X(1) &= 1 \\ X(2) &= 2 \\ X(3) &= 3 \\ X(4) &= 4 \\ X(5) &= 5 \\ X(6) &= 6 \end{aligned}$$

$$p_X(k) = \frac{1}{6} \quad k = 1, \dots, 6$$

Q: What's the probability that X is even?

$$\rightarrow: \mathbb{P}[X \text{ is even}] = \mathbb{P}[X \in \{2, 4, 6\}] = \frac{1}{2}$$

Let Y be the random variable defined as:

$$Y = \begin{cases} 1 & \text{if the result is prime} \\ 0 & \text{otherwise} \end{cases}$$

$$\begin{aligned} Y(1) &= 0 \\ Y(2) &= 1 \\ Y(3) &= 1 \\ Y(4) &= 0 \\ Y(5) &= 1 \\ Y(6) &= 0 \end{aligned}$$


$$\text{Support}(Y) = \text{Range}(Y) = \{0, 1\}$$

$$p_Y(0) = \frac{1}{2}$$

$$p_Y(1) = \frac{1}{2}$$



Functions of random variables.



$$g(X(\omega)) = g(X)(\omega)$$

A function of a random variable is (again) a random variable.

We assign it a new "Label", e.g.,

$$Y = g(X)$$

We can look @ its distribution 😊